

Name: Key

Student ID: _____

No Calculators.**Show all work and justifications to receive full credit.**

1. (4 pts) Find the angle between the two planes $7x + 7y + 7z = 7$ and $3x - 10y + 5z = -8$. You do not need to simplify your answer.

To find the angle between two planes is equivalent to finding the angle between the normals to the plane.

$$\langle 7, 7, 7 \rangle \text{ or } \langle 1, 1, 1 \rangle = n_1 \text{ is the normal to plane } 7x + 7y + 7z = 7 \quad \& \quad \langle 3, -10, 5 \rangle = n_2 \text{ is the normal to plane } 3x - 10y + 5z = -8$$

Thus $\cos \theta = \frac{n_1 \cdot n_2}{\|n_1\| \|n_2\|}$

$$= \frac{-2}{\sqrt{3^2 + 7^2}} = \frac{-2}{\sqrt{50}} \Rightarrow \theta = \cos^{-1} \left(\frac{-2}{\sqrt{50}} \right)$$

2. (4 pts) Find the domain and range of $f(x, y) = \frac{1}{\sqrt{7-x^2-y^2}}$.

The domain (f) = $\{(x, y) \mid 7-x^2-y^2 > 0\} = \{(x, y) \mid x^2+y^2 < 7\}$
 = points in the circle of radius $\sqrt{7}$ centered at the origin

range (f) = $[\frac{1}{\sqrt{7}}, \infty)$ since the largest $7-x^2-y^2$ is 7 & smallest is close to 0.

3. (2 pts) Describe the level curve $1 = z = f(x, y)$ where $f(x, y)$ is defined in problem 2.

$$1 = \frac{1}{\sqrt{7-x^2-y^2}} \Rightarrow \sqrt{7-x^2-y^2} = 1 \Rightarrow 7-x^2-y^2 = 1$$

$\Rightarrow x^2+y^2 = 6$ which is a circle of radius $\sqrt{6}$ centered at the origin.